

Dual fields and E_{11}

Fabio Riccioni and Peter West
Department of Mathematics
King's College, London WC2R 2LS, UK

Abstract

We show that the adjoint representation of E_{11} contains generators corresponding to the infinite possible dual descriptions of the bosonic on-shell degrees of freedom of eleven dimensional supergravity. We also give an interpretation for the fields corresponding to many of the other generators in the adjoint representation.

A few years ago it was conjectured [1] that a rank eleven Kac-Moody algebra, which was called E_{11} , was a symmetry of M theory. The non-linear realisation of this algebra at its lowest levels was shown to contain the fields of eleven dimensional supergravity and to have the equations of motion of this theory [1] when one made use of the earlier results of reference [2]. The physical field content is extracted by decomposing the adjoint representation of E_{11} into the representations of its A_{10} , or $Sl(11)$, sub-algebra which is associated in the non-linear realisation with eleven dimensional gravity. Non-linear realisations of E_{11} can also be used to describe ten dimensional theories, but in this case the adjoint representation is decomposed into representations of A_9 , or $Sl(10)$, associated with ten dimensional gravity. There are only two such A_9 subalgebras and the two different choices were found to lead to non-linear realisations which at low levels are the IIA and IIB supergravity theories in references [1] and [3] respectively. It is striking to examine tables of the generators [4] listed in terms of increasing level and see how the generators associated with the field content of the IIA and IIB supergravity theories occupy precisely all the lower levels before an infinite sea of generators whose physical significance was unknown at the time the tables of reference [4] were constructed.

Amongst the set of E_{11} generators appropriate to the IIA theory is one with nine antisymmetrised indices which in the non-linear realisation leads to a field with the same index structure [4]. This nine form generator occurs in the table of IIA generators at a place which is amongst the generators associated with the fields of the IIA supergravity theory. A non-trivial value for this field is known to lead to the massive IIA supergravity theory and as a result it was realised [4] that E_{11} can incorporate the massive IIA theory. However, there is a one to one correspondence between the fields of the non-linear realisations of E_{11} appropriate to the eleven dimensional, IIA and IIB theories and using this correspondence one finds that the nine form of the IIA theory corresponds to a field with the index structure $A^{a,b,c_1\dots c_{10}}$ which occurs at a level which is beyond those of the supergravity fields of eleven dimensional supergravity theory [5]. As such E_{11} provides for an eleven dimensional origin of the massive IIA theory which involves one of the higher level fields and in this way the physical interpretation of at least one of the higher level fields in the eleven dimensional E_{11} theory became apparent.

The non-linear realisation of E_{11} appropriate to the IIB theory was observed to contain a number of eight form and ten form fields at levels just above the usual fields of supergravity [4]. The IIB supergravity theory was originally constructed by closing the local supersymmetry algebra [6]. One can add to the standard supergravity fields a $SL(2,R)$ doublet of six forms, dual to the two forms, and a triplet of eight forms dual to the scalars, whose 9-form field strengths satisfy a constraint so that only two fields actually propagate [9], but in an unexpected recent development [7] it was shown that one could also add a very specific set of ten forms to the theory and still close the supersymmetry algebra. Remarkably, these fields were in precise correspondence with those already found in E_{11} [4]. Furthermore, the exact coefficients of their gauge transformations deduced from supersymmetry closure [7] were found [8] to agree with their derivation from the E_{11} algebra. The ten form fields, although non-propagating, have an important role since they are associated with spacetime-filling branes. In particular, one of the 10-forms in the IIB algebra is associated to D9-branes, that take part in the orientifold projection which gives rise

to the type-I theory from the IIB theory [10]. Thus a few more of the higher level fields were found to have a physical meaning. The non-linear realisation of E_{11} appropriate to IIA theory was also observed to contain some ten forms [4] and these were recently found [11] to be in precise agreement with the ten forms one could add and still have a closing supersymmetry algebra in the IIA theory.

In this paper we will consider the generators of the adjoint representation of E_{11} classified according to the A_{10} sub-algebra corresponding to an eleven dimensional gravity subsector. We will find all generators that have no blocks of A_{10} indices containing ten antisymmetrised indices and whose indices sum to $3l$ where l is the level of the generator. The lowest level such generators have the index structure R^{abc} , $R^{a_1\dots a_6}$, $R^{a_1\dots a_8,b}$ and correspond to the three and six form gauge fields of the supergravity theory and the dual graviton field. Although there could in principle be a very large number of generators in the class being considered we will show that it only contains the above generators and the above generators decorated by blocks each of which contain nine indices. For example, for the three form this means the index structure $R^{abc,d_1\dots d_9,e_1\dots e_9}$ when we have only two blocks of nine indices. The fields corresponding to these generators are just all possible dual formulations of the bosonic on-shell degrees of freedom of eleven dimensional supergravity. We interpret this result to mean that this sector of the adjoint representation of E_{11} encodes the most general possible duality symmetries of bosonic eleven dimensional supergravity in a manifest manner. As a result, we will find a physical interpretation for an infinite number of E_{11} generators.

Very little is known about generalised Kac-Moody algebras, indeed, for no such algebra is there even a listing of the generators that are present in the algebra. However, Kac-Moody algebras which possess at least one node whose deletion leads to a set of Dynkin diagrams each of which corresponds to one of the finite dimensional semi-simple Lie algebras are more amenable to analysis [12]. In particular, one can analyse the Kac-Moody algebra in terms of the finite dimensional sub-algebras that appears after this deletion. The Dynkin diagram of E_{11} is given by

$$\begin{array}{cccccccccccc}
 & & & & & & & & & & 0 & 11 \\
 & & & & & & & & & & | & \\
 0 & - & 0 & - & 0 & - & 0 & - & 0 & - & 0 & - & 0 & - & 0 & - & 0 \\
 1 & & 2 & & 3 & & 4 & & 5 & & 6 & & 7 & & 8 & & 9 & & 10
 \end{array}$$

In the case of E_{11} , if one deletes the node labeled eleven then the remaining algebra is A_{10} or $SL(11)$. We call this A_{10} section of the Dynkin diagram the gravity line as it gives rise in the non-linear realisation to eleven dimensional gravity sector of the theory. It will prove useful to recall how one analyses the generators of the adjoint representation E_{11} in terms of representations of A_{10} [12,13,14]. The simple roots of E_{11} are the simple roots α_i , $i = 1, \dots, 10$ of A_{10} as well as the simple root corresponding to node eleven which we call α_c which is given by

$$\alpha_c = x - \lambda_8 \quad (1)$$

Here x is orthogonal to the roots α_i , $i = 1, \dots, 10$ of A_{10} and λ_i , $i = 1, \dots, 10$ are the fundamental weights of A_{10} . As $\alpha_c \cdot \alpha_c = 2$ and $\lambda_8 \cdot \lambda_8 = \frac{8.3}{11}$, we find that $x^2 = -\frac{2}{11}$. Any

root α of E_{11} can be expressed in terms of its simple roots of E_{11} and, using equation (1), can be written as follows

$$\alpha = \sum_{i=1}^{10} n_i \alpha_i + l \alpha_c = lx - \Lambda \quad (2)$$

where

$$\Lambda = l \lambda_8 - \sum_{i=1}^{10} n_i \alpha_i \quad (3)$$

is in the weight space of A_{10} .

We call l the level of the root and the strategy [12,13,14] is to analyse the generators in the adjoint representation of E_{11} level by level in terms of the representations of A_{10} . When constructing the E_{11} algebra in terms of multiple commutators using the Serre relation, the level is just the number of times the generator R^{abc} occurs in the commutators.

If the adjoint representation of E_{11} contains the representation of A_{10} at level l with Dynkin indices p_j , then we must find that some root of E_{11} contains an A_{10} weight of the form $\Lambda = \sum_j p_j \lambda_j$ where p_j are positive integers. As such, one finds that

$$\sum_j p_j \lambda_j = l \lambda_8 - \sum_j n_j \alpha_j \quad (4)$$

Taking the scalar product with λ_i , and using the equation

$$(A_{ik})^{-1} = (\lambda_i, \lambda_k) \quad (5)$$

valid for any simply laced finite dimensional semi-simple Lie algebra, implies that

$$\sum_j p_j A_{ji}^{-1} = l A_{8i}^{-1} - n_i \quad (6)$$

where A_{ji}^{-1} is the inverse Cartan matrix of A_{10} . After some thought one can see that this must also occur for p_j , l and n_i are all positive integers. As such equation (6) places strong restrictions on the possible p_j , and so the representations of A_{10} , that can occur at a given level. This observations relies on the fact that the inverse Cartan matrix is positive definite for any finite dimensional semi-simple Lie algebra.

Any Kac-Moody algebra with symmetric Cartan matrix has its roots bounded by $\alpha^2 \leq 2, 0, -2 \dots$ [15]. Consequently, using equations (2) and (5) we find that

$$\alpha^2 = l^2 x^2 + \sum_{ij} p_i (A_{ij})^{-1} p_j \leq 2, 0, -2, \dots \quad (7)$$

Since the first term is fixed for a given l and the second term is positive definite this also places constraints on the possible values of p_i .

It will be useful to recall the generators of E_{11} that occur for the first three levels [1]. At level zero we have the generators K^a_b of A_{10} , at level one

$$R^{abc}, \quad l = 1, \quad p_8 = 1 \quad \alpha_A = (0, 0, 0, 0, 0, 0, 0, 0, 0, 1) \quad (8)$$

at level two

$$R^{a_1 \dots a_6}, \quad l = 2, \quad p_5 = 1 \quad \alpha_B = (0, 0, 0, 0, 0, 1, 2, 3, 2, 1, 2) \quad (9)$$

at level three

$$R^{a_1 \dots a_8, b}, \quad l = 3, \quad p_{10} = 1, p_3 = 1 \quad \alpha_C = (0, 0, 0, 1, 2, 3, 4, 5, 3, 1, 3) \quad (10)$$

The vectors shown above are the E_{11} roots, the entries being the integers of equation (2). At level three we also find

$$R^{a_1 \dots a_9}, \quad p_2 = 1 \quad \alpha_D = (0, 0, 1, 2, 3, 4, 5, 6, 4, 2, 3), \quad (11)$$

but this generator has multiplicity zero and so does not actually occur in the E_{11} algebra. All other p_j 's not mentioned in the above expressions are zero. In deriving these results one uses that the inverse Cartan matrix of A_{n-1} , or $SL(n)$, is given by

$$(A_{jk})^{-1} = \begin{cases} \frac{j(n-k)}{n}, & j \leq k \\ \frac{k(n-j)}{n}, & j \geq k \end{cases} \quad (12)$$

We now show that the E_{11} generators at level l when decomposed into representations of A_{10} can be written in such a way that they always have $3l$ indices. This includes level zero if one counts upper indices with a plus sign and lower indices with a minus sign. Clearly, this counting is not respected when changing indices using the epsilon symbol. Like all Kac-Moody algebras, E_{11} is just the multiple commutators of its Chevalley generators, subject to the Serre relation. The multiple commutators of the level zero Chevalley generators lead to $SL(11)$, or A_{10} that is the generators K^a_b . While the multiple commutators of these, together with one of the one level one Chevalley generator, R^{91011} lead to the generators R^{abc} . The rest of the positive root part of the algebra is then found from multiple commutators of R^{abc} . As a result, if one never uses epsilons on the right hand side of the commutators one finds generators that have $3l$ $SL(11)$ indices. The same applies to the construction of all the negative root generators which can also be written with $3l$ indices, but now the levels are negative. This is consistent with the assignment of a lower index with -1 contribution.

The generators that emerge from the above analysis are representations of A_{10} and so are labeled by their associated Dynkin indices p_j . In our conventions a generator with $p_j = 1$ has $11 - j$ antisymmetrised indices. To each non-vanishing Dynkin index, p_j the generator possess p_j blocks of antisymmetric indices. The blocks of antisymmetrised indices are in general subject to symmetrisation relations.

We now divide all the generators into two classes, those that have at least one block of indices that contain ten or eleven antisymmetrised indices and those generators that do not contain any such blocks. We refer to the latter generators as **dual** generators for a reason that will become apparent. We note that a block of eleven antisymmetrised indices is equivalent to a scalar and so does not correspond to any Dynkin index. However, one can count the number of such blocks as it is just determined by the mismatch between the sum of the indices that arise from the blocks associated with the Dynkin indices p_j and $3l$.

The dual generators have, by definition, no such rank eleven index blocks and also have the Dynkin index $p_1 = 0$, as they also have no blocks of indices with ten antisymmetrised indices. As such, dual generators have $p_1 = 0$ and sum of all the indices must be $3l$;

$$\sum_{j=2}^{10} p_j(11-j) = 3l \quad (13)$$

Let us give some more examples. At level four in the table of E_{11} generators we find a generators with Dynkin indices $p_2 = 1$ and $p_8 = 1$, all other Dynkin indices being zero. This has the index structure $R^{a_1 a_2 a_3, b_1 \dots b_9}$, it has the required $3.4 = 12$ indices and so is a dual generator. Also at level four we find a generator with $p_{10} = 1$, all other Dynkin indices being zero. This must be of the form $R^{a_1, b_1 \dots b_{11}}$ in order to have the required 12 indices and so it is not a dual generator.

We now use equations (6) and (7) to see what constraints they place on the possible dual generators. We begin with the condition on the length of the root of equation (7). Substituting for l from equation (13) we find that equation (7) considerably simplifies and can be written as

$$\alpha^2 = \frac{1}{9} \sum_j (11-j)(j-2)p_j^2 + \frac{2}{9} \sum_{i < j} (11-j)(i-2)p_i p_j = 2, 0, -2, \dots \quad (14)$$

We observe that the left hand side is positive definite and so the root α can only have length squared 2 or 0. Furthermore, we note that the equation does not contain p_2 and so places no constraint on it. It is straightforward to show that the only solutions are

$$\text{solution A,} \quad p_2 = u, p_8 = 1, \quad \text{at } l = 1 + 3u, \quad (15)$$

$$\text{solution B,} \quad p_2 = u, p_5 = 1, \quad \text{at } l = 2 + 3u, \quad (16)$$

$$\text{solution C,} \quad p_2 = u, p_3 = 1 = p_{10}, \quad \text{at } l = 3 + 3u, \quad (17)$$

$$\text{solution D,} \quad p_2 = u + 1, , \quad \text{at } l = 3 + 3u, \quad (18)$$

All these solutions automatically satisfy equation (6) and one finds that the E_{11} roots they correspond to are given by

$$\alpha_I(u) = \alpha_I + u\alpha_D \quad (19)$$

where the label $I = A, B, C, D$ corresponding to the solutions and α_I are given in equations (8-11).

We first discuss the solutions A,B,C which all have $\alpha^2 = 2$. These contain just one solution at every level. The solutions for $u = 0$ are just the three form R^{abc} at level one, the six form $R^{a_1 \dots a_6}$ at level two and the dual graviton generator $R^{a_1 \dots a_8, b}$ at level three given in equations (8), (9) and (10). This must be the case as all these generators are dual generators. At higher levels this pattern is repeated, but one finds instead that the generators acquire u blocks of nine indices. Solution D had $\alpha^2 = 0$ and for $u = 0$ it is

just the nine form at level three of equation (11). At all higher levels spaced by three one finds in solution D generators with $u + 1$ blocks of nine indices. In fact the level three, i.e. $u = 0$, nine form generator $R^{a_1 \dots a_9}$, has multiplicity zero and examining the E_{11} tables it would appear that all the other generators in solution D also have multiplicity zero up to level nine and one can expect this to be true at all the levels. This means that although these generators solve all the constraints of equations (6) and (7) they do not actually occur in the algebra. The constraints of equations (6) and (7) are a necessary, but not a sufficient condition. What generators actually occur in the algebra are determined by multiple commutators of the Chevalley relations subject to the Serre relation and this is more restrictive than the above conditions. In contrast, looking at the tables of reference [16,19] we see that the generators for solutions A, B and C all have multiplicity one up to level ten and so do occur in the algebra. We can expect that they have multiplicity one at all levels.

Although equations (6) and (7) can be used to systematically search for a generator with fixed Dynkin indices at all levels, there is usually little hope of successfully carrying out such a calculation unless the Dynkin indices are specified. However, this was not the case for the dual generators indicating that they are preferred objects in the E_{11} algebra.

The massless states of the eleven dimensional Poincare algebra are classified in terms of representations of its little group $SO(9)$. If we take the indices on the generators in the solutions A, B and C in equations (15-17) to only take the values $1, \dots, 9$, the blocks of nine indices are equivalent to no indices at all and so we find only the three rank, six rank and the $R^{j_i \dots j_8, i}$ irreducible representations of $SO(9)$. Of course the rank three and rank six generators are the same representations of $SO(9)$ as is apparent by invariant epsilon symbol. The same is true for the graviton contained in the level zero sector and the $R^{j_i \dots j_8, i}$ generator at level three as the latter is just equivalent to the generator $R_j{}^i$. In fact the generator $R^{j_i \dots j_8, i}$ obeys the constraint $R^{[j_1 \dots j_8, j_9]} = 0$ and so the equivalent generator obeys $R_i{}^i = 0$. The fields corresponding to the generators of the solutions A, B and C are just all possible dual descriptions of the bosonic degrees of freedom of the eleven dimensional supergravity theory. The meaning of this result is that the sector of the E_{11} non-linear realisation associated with the dual generators encodes all possible duality symmetries in a manifest way. We note that all the non-dual fields vanish when we restrict their indices to only take the values $1, \dots, 9$ and so they lead to no representations of the little group $SO(9)$. This suggests that the non-dual generators lead to no propagating degrees of freedom.

The roots of equation (19) for the generators in the solutions of equations (15-17) are consistent with the conjecture that all these generators are found taking multiple commutators of the generator $R^{a_1 \dots a_9}$ of equation (18) with the three generators of equations (8-10).

We now apply similar considerations to the E_{11} theory appropriate to the IIB theory. The Dynkin diagram of E_{11} with the A_9 subgroup associated with the gravity sector of

the IIB theory shown as the horizontal line (the so called gravity line) [3].

$$\begin{array}{ccccccccccccccc}
 & & & & & & & & & & & & 0 & 10 \\
 & & & & & & & & & & & & | & \\
 & & & & & & & & & & & & 0 & 9 \\
 & & & & & & & & & & & & | & \\
 0 & - & 0 & - & 0 & - & 0 & - & 0 & - & 0 & - & 0 & - & 0 \\
 1 & & 2 & & 3 & & 4 & & 5 & & 6 & & 7 & & 8 & - & 11
 \end{array}$$

It is immediately apparent that the IIB theory has an $SL(2, \mathbb{R})$ invariance which arises from the node labeled ten as this is not connected to the gravity line and so corresponds to an internal symmetry. For this theory we delete the node labeled 9 and then the algebra splits into $A_9 \otimes A_1$. The simple roots of E_{11} are the simple α_i , $i = 1, \dots, 8, 11$ of A_9 , the root α of A_1 and the root corresponding to node 9 which we call α_c which is given by

$$\alpha_c = x - \mu - \lambda_8 \quad (20)$$

where λ_i , $i = 1, \dots, 8, 11$ are the fundamental weights of A_9 and μ is the fundamental weight of A_1 . We note that the last node in the gravity line is node 11 and we keep this labeling for the A_9 Dynkin diagram. As $\alpha_c \cdot \alpha_c = 2$ we find that $x^2 = -\frac{1}{10}$. Any root α of E_{11} can be expressed in terms of the simple roots of E_{11} as follows

$$\alpha = \sum_{i=1}^{8,11} n_i \alpha_i + l \alpha_c + m \alpha = l x - \Lambda^{A_9} - \Lambda^{A_1} \quad (21)$$

where

$$\Lambda^{A_9} = l \lambda_8 - \sum_{i=1}^{8,11} n_i \alpha_i, \quad \text{and} \quad \Lambda^{A_1} = l \mu - m \alpha \quad (22)$$

We note that Λ^{A_9} and Λ^{A_1} are in the weight spaces of A_9 and A_1 respectively. A necessary condition for the adjoint representation of E_{11} to contain the representation of $A_9 \otimes A_1$, with Dynkin indices p_j and q for each factor respectively, is that

$$\Lambda^{A_9} = l \lambda_8 - \sum_{i=1}^{8,11} n_i \alpha_i = \sum_j p_j \lambda_j, \quad (23)$$

and

$$\Lambda^{A_1} = l \mu - m \alpha = q \mu \quad (24)$$

Where l, n_i, p_i and m, q are all positive integers or zero. Taking the scalar product with the fundamental weights, these two equations respectively become

$$\sum_j p_j A_{ji}^{-1} = l A_{8i}^{-1} - n_i \quad (25)$$

and

$$l - q = 2m \quad (26)$$

where now A_{ji}^{-1} is the inverse Cartan matrix of A_9 . The IIB decomposition given in reference [4] has two levels l_1 and l_2 corresponding to the deletion of the nodes ten and nine. The decomposition studied here has the advantage that it automatically works with complete representations of $SL(2, R)$. For comparison we have $l_2 = l$ while $l_1 = m = \frac{1}{2}(l - q)$

The condition on the length of the roots is given by

$$\alpha^2 = l^2 x^2 + \sum_{ij} p_i (A_{ij})^{-1} p_j + \frac{q^2}{2} \leq 2, 0, -2, \dots \quad (27)$$

We now divide the generators of the IIB theory into two classes in a way that is similar to that done above for the eleven dimensional theory. The generators of the IIB theory can all be written in such a way that they carry $2l$ indices. This is essentially because the Chevalley generators of the IIB theory are those associated with the gravity line nodes one to eight and node eleven, whose multiple commutators lead to the generators K^a_b of A_9 , the Chevalley generator of node ten which is associated with $SL(2, R)$, whose multiple commutators introduce no space-time indices and finally the Chevalley generator of node nine which is contained in one of the two generators of the form R^{ab} and has level one. It is the multiple commutators of this latter generators that lead to the addition of two space-time indices at every level. We now call dual generators those that have no blocks of antisymmetrised indices of rank nine or ten. As a result, the dual generators have $p_1 = 0$ and

$$\sum_{j=2}^{8,11} p_j (10 - j) = 2l \quad (28)$$

Using equation (28) we find the length condition on the root of equation (27) leads to the constraint

$$\alpha^2 = \frac{1}{8} \sum_j (10 - j)(j - 2) p_j^2 + \frac{1}{4} \sum_{i < j} (10 - j)(i - 2) p_i p_j + \frac{q^2}{2} = 2, 0, -2, \dots \quad (29)$$

We see that the left hand side is positive definite and so the only non-trivial solutions have $\alpha^2 = 2$. We also observe that the Dynkin label p_2 does not appear and so it not subject to any constraint. Furthermore q can only take the values $q = 0, 1, 2$. If we have $q = 2$ then all the $p_j = 0$ except p_2 which can take any value. It is easy to see that the only solutions, which also satisfy equations (25) and (26), are

$$\text{solution A, } R^{a_1 a_2}, \quad p_2 = u, p_8 = 1, \quad q = 1, \quad \text{at } l = 1 + 4u, \quad (30)$$

$$\text{solution B, } R^{a_1 \dots a_4}, \quad p_2 = u, p_6 = 1, \quad q = 0 \quad \text{at } l = 2 + 4u, \quad (31)$$

$$\text{solution C, } R^{a_1 \dots a_6}, \quad p_2 = u, p_4 = 1, \quad q = 1 \quad \text{at } l = 3 + 4u, \quad (32)$$

$$\text{solution D, } R^{a_1 \dots a_7, b}, \quad p_2 = u, p_3 = 1 = p_9, \quad q = 0 \quad \text{at } l = 4 + 4u, \quad (33)$$

$$\text{solution E, } R^{a_1 \dots a_8}, \quad p_2 = u + 1, \quad q = 2 \quad \text{at } l = 4 + 4u, \quad (34)$$

$$\text{solution F, } R^{a_1 \dots a_8}, \quad p_2 = u + 1, \quad q = 0 \quad \text{at } l = 4 + 4u, \quad (35)$$

The above only lists the highest weight components of each $SL(2, R)$ multiplet and one can obtain all members of the multiplet by acting with the Chevalley generators E_{10} and F_{10} . The axion is not displayed as it is part of the $SL(2, R)$ multiplet which contains the dilaton as its highest weight component. We notice that for $u = 0$ we have precisely the generators corresponding to the content of the IIB supergravity theory if we include the dual graviton and the eight form generators. The above solutions all have $\alpha^2 = 2$ except for solution F which has $\alpha^2 = 0$. The general solution above has generators which have the indices of the $u = 0$ generators decorated by any number of blocks of eight indices. The roots are found from equations (25) and (26). The simplest way to deduce them is to first show that if one has a solution with certain set of Dynkin indices p_j , then there exists another solution with the same Dynkin indices except with p_2 replaced by $p_2 + u$ for u a positive integer. The resulting E_{11} root α then changes to $\alpha + u\alpha_F$. Hence, given the above solutions for $u = 0$ we generate all the others and they have the roots

$$\alpha_I(u) = \alpha_I + u\alpha_F \quad (36)$$

where I stands for A,B,C,D,E and F and the roots α_I are given by

$$\begin{aligned} \alpha_A &= (0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0), & \alpha_B &= (0, 0, 0, 0, 0, 0, 1, 2, 2, 1, 1), \\ \alpha_C &= (0, 0, 0, 0, 1, 2, 3, 4, 3, 1, 2), & \alpha_D &= (0, 0, 0, 1, 2, 3, 4, 5, 4, 2, 2), \\ \alpha_E &= (0, 0, 1, 2, 3, 4, 5, 6, 4, 1, 3), & \alpha_F &= (0, 0, 1, 2, 3, 4, 5, 6, 4, 2, 3) \end{aligned} \quad (37)$$

The little group in ten dimensions is $SO(8)$. The above solutions have a similar interpretations as the ones in eleven dimensions. They correspond to all possible dual descriptions of the on-shell bosonic degrees of freedom of the IIB supergravity theory and so one expects that the non-linear realisation corresponding to the IIB theory manifestly displays all possible duality symmetries corresponding to the on-shell degrees of IIB supergravity. As for the eleven dimensional case, the non-dual fields do not contribute any non-trivial little group representations. This completes our discussion of the IIB case.

We observed that for the eleven dimensional case the dual generators were likely to be result from taking repeated commutators of the generator at level three which had $p_2 = 1, l = 3, \alpha^2 = 0$ and multiplicity zero with the three lowest level generators. In fact such a result applies not only to the dual generators, but to all the E_{11} generators as we now explain. Given two solutions with Dynkin indices and levels given by $(p^{(1)}, l^{(1)})$ and $(p^{(2)}, l^{(2)})$ to the weight matching condition of equation (6), then we find that $(p^{(1)} + p^{(2)}, l^{(1)} + l^{(2)})$ is also a solution as this equation is linear in its Dynkin indices and level. This is not in general true for the root length condition of equation (7). However, if we take $(p^{(2)}, l^{(2)})$ to be the generator of equation (18) with $u = 0$, i.e. $(p_2^{(2)} = 1, l^{(2)} = 3)$, then we do find a solution of equation (7) of the form $(p^{(1)} + p^{(2)}, l^{(1)} + l^{(2)})$. Indeed, if we denote the root of the latter by β then it has a root length squared which is given by

$$(\beta)^2 = \alpha_1^2 - \frac{2 \cdot 2}{11} l^{(1)} l^{(2)} + \sum_{i,j} p_j^{(1)} A_{ji}^{-1} p_j^{(2)} = \alpha_1^2 - 2n_2^{(1)} \quad (38)$$

which is always in the allowed range. In going between the two equations we have used equation (6) for the first solution. Thus even though the generator with $p_2 = 1, l = 3$ does not actually appear in the E_{11} algebra its repeated commutator with any generator in the E_{11} algebra should result in another generator in the algebra. Examining the table of E_{11} generators [16,19] one does indeed find generators which are genuine members of the E_{11} algebra with the correct structure to have arisen from such commutators. Such generators will have the same $SO(9)$ little group representations as the original generator before any commutators and as such it would seem that all fields in the non-linear realisation of E_{11} come in all possible dual formulations.

The E_{11} root corresponding to the generator $R^{a_1 \dots a_9}$ with $p_2 = 1, l = 3$ is of the form $k \equiv \alpha_3 + \theta$ where θ is the highest root of E_8 and α_3 is the root whose addition corresponds to changing E_8 to the affine algebra E_9 . As such, k is associated with the null root of the affine E_9 subalgebra. We note that in any affine algebra the null root k is responsible for the infinite number of generators in the algebra as if α is a root of length squared two found in the underlying finite dimensional algebra then the roots of the algebra are of the form $\alpha + nk$. The infinite number of generators in the affine algebra can be packaged up into $T^a(z) = T_n^a z^n$ where z parameterises the S^1 of the circle or loop. Given the above results we conclude that we can package all the generators of E_{11} in a similar manner by considering them as coefficients of a polynomial in the variable $z^{a_1 \dots a_9}$. Even once this packaging has been carried out the adjoint representation contains an infinite number of generators even though they are functions of $z^{a_1 \dots a_9}$. However, the dual generators can all be encoded by taking the generators of the adjoint representation of E_8 and letting them depend on $z^{a_1 \dots a_9}$. Could it be that E_{11} can be formulated as a mapping from a three dimensional surface into the E_8 algebra and could the same apply to any very extended algebra G^{+++} ?

Similar results to those discussed just above hold for the IIB theory using the generator of equation (35).

Given any finite dimensional semi-simple Lie algebra G one can form a unique rank three larger Kac-Moody algebra G^{+++} based upon it [11] and consider the non-linear realisation formed from it [1,17,18]. For example, if $G = E_8$ then $E_8^{+++} = E_{11}$. The low level dynamics of these non-linear realisations are many of the theories of interest to theoretical physicists. For example, the non-linear realisations of A_{D-3}^{+++} and D_{D-2}^{+++} are gravity [1] and a generalisation of the effective action of the bosonic string [17] in D space-time dimensions respectively. Examining the tables of references [4] and [19] one finds that the story of the dual generators found in the context of E_{11} also holds for these theories. Namely, the dual generators are the original low level generators associated with the physical fields of the theory as well as these generators when supplemented by blocks of $D - 2$ indices. Furthermore there exists in each case generators of length squared zero whose multiple commutators with the very lowest dual generators are likely to lead to all dual generators. This is particularly apparent when looking at the table of generators of D_8^{+++} and B_3^{+++} of reference [4] and A_1^{+++} of reference [19].

Let us summarise the results in this paper, we have considered the adjoint representation of E_{11} when decomposed with respect to the A_{10} sub-algebra corresponding to the gravity sector of an eleven dimensional theory. We have explained that all the generators

of E_{11} can be written with $3l$ indices where l is the level of the generator. We then found all generators of E_{11} that do not have any blocks of ten or eleven antisymmetrised indices. There is only one such generator at every level and they correspond to all possible dual formulations of the on-shell bosonic degrees of freedom of the eleven dimensional supergravity theory. Thus we have found a physical meaning for an infinite number of E_{11} generators.

The non-dual generators, when written with indices as described in this paper, lead to no degrees of freedom for the little group and one may suspect they do not correspond to propagating degrees of freedom. One word of caution is required in relation to this last statement as it depends on writing all generators with $3l$ indices, however, one can change the number of indices with the epsilon symbol and only the dynamics can determine which fields actually propagate. Only once one has the dynamics can one determine what the actual propagating degrees of freedom are.

The corresponding generators in the IIB theory follow a similar pattern and interpretation and it is likely that the results hold for any non-linear realisation based on any very extend algebra G^{+++} .

Given the results in this, and previous papers on E_{11} , there is now considerable evidence that E_{11} is closely associated with the maximal supergravity theories. Indeed, it would seem increasingly likely that the original conjecture of reference [1] that E_{11} is a symmetry of a suitably decorated formulation of eleven dimensional supergravity is true. Since the maximal ten dimensional supergravity theories are the complete low energy effective actions for the corresponding superstrings this result also encourages the belief that the underlying theory has an E_{11} symmetry, albeit of a discrete kind.

The dual generators are only a small part of the infinite number of generators of the adjoint representation of E_{11} . However, as explained in the introduction a number of the non-dual generators, although non-propagating, have a physical meaning. For example, the level four generator $R^{a,b,c_1\dots c_{10}}$ of the eleven dimensional theory allows for the emergence of the massive IIA supergravity theory upon reduction to ten dimensions. One can expect that this is a general phenomenon, namely the gauged, sometimes called massive, supergravities that occur in lower dimensions can be thought of as arising from fields beyond the supergravity approximation in the eleven dimensional E_{11} non-linear realisation. Indeed, it would be interesting to see if a precise correspondence can be made between such fields and the gauged supergravities in say three dimensions. Thus one expects to find an interpretation for many of the non-dual generators and these results encourage the belief that all the fields in the non-linear realisation of E_{11} have physical meaning.

In this paper we have only shown that the field content of the non-linear realisations corresponds to all possible dual fields. However, it would be good to show that the dynamics of the non-linear realisation actually leads to a formulation with all possible dual field incorporated in the correct way. It is possible that the dual generators and the generator with $u = 0$ of equation (18) form a closed subalgebra which is a truncation of the E_{11} with the generators of equation (11) added. If this were the case there would be a manifest description of the duality symmetries of the on-shell degrees of freedom without the non-dual fields.

Acknowledgments Peter West would like to thank Rafael Montemayor for important

discussions at an early stage of this work and hospitality at the centre for physics at Bariloche in January 2005 and at the physics department of the University of Canterbury at Christchurch, New Zealand in November 2006. He would also like to thank Andrew Pressley for discussions on group theory. Fabio Riccioni would like to thank E. Bergshoeff for discussions. The research of Peter West was supported by a PPARC senior fellowship PPA/Y/S/2002/001/44. The work of both authors is also supported by a PPARC rolling grant PP/C5071745/1 and the EU Marie Curie, research training network grant HPRN-CT-2000-00122.

References

- [1] P. West, *E₁₁ and M Theory*, Class. Quant. Grav. **18** (2001) 4443, hep-th/010408.
- [2] P. C. West, *Hidden superconformal symmetry in M theory*, JHEP **08** (2000) 007, hep-th/0005270
- [3] I. Schnakenburg and P. West, *Kac-Moody Symmetries of IIB supergravity*, Phys. Lett. **B 517** (2001) 137-145, hep-th/0107181
- [4] A. Kleinschmidt, I. Schnakenburg and P. West, *Very-extended Kac-Moody algebras and their interpretation at low levels*, hep-th/0309198
- [5] P. West, *The IIA, IIB and eleven dimensional theories and their common E₁₁ origin*, hep-th/0402140.
- [6] J. Schwarz and P. West *Symmetries and Transformations of chiral N=2, D=10 supergravity*, Phys. Lett. **B126** (1983), 301; J. Schwarz, *Covariant field equations of chiral N=2 D=10 supergravity*, Nucl. Phys. **B226** (1983), 269; P. Howe and P. West, *The complete N=2, d=10 supergravity*, Nucl. Phys. **B238** (1984) 181.
- [7] E. Bergshoeff, Mess de Roo, S. Kerstan and F. Riccioni, *IIB Supergravity Revisited*, hep-th/0506013.
- [8] P. West, *E₁₁, ten forms and supergravity*, JHEP 0603 (2006) 072, hep-th/0511153.
- [9] P. Meessen and T. Ortin, *An SL(2,Z) multiplet of nine -dimensional type II supergravity theories*, Nucl. Phys. **B 541** (1999) 195, hep-th/9806120; G. Dall'Agata, K. Lechner and M. Tonin, *D=10, N=IIB supergravity; Lorentz-invariant actions and duality*, JHEP **9807** (1998) 017, hep-th/9806140;
- [10] A. Sagnotti, *Open strings and their symmetry groups*, hep-th/0208020.
- [11] Eric A. Bergshoeff, Mees de Roo, Sven F. Kerstan, T. Ortin, Fabio Riccioni, *IIA Ten-forms and the Gauge Algebras of Maximal Supergravity Theories*, JHEP 0607 (2006) 018, hep-th/0602280.
- [12] M. R. Gaberdiel, D. I. Olive and P. West, *A class of Lorentzian Kac-Moody algebras*, Nucl. Phys. **B 645** (2002) 403-437, hep-th/0205068.
- [13] T. Damour, M. Henneaux and H. Nicolai, *E₁₀ and a "small tension expansion" of M-theory*, Phys. Rev. Lett. **89** (2002) 221601, hep-th/0207267
- [14] P. West, *Very Extended E₈ and A₈ at low levels*, Gravity and Supergravity, Class. Quant. Grav. **20** (2003) 2393-2406, hep-th/0212291.
- [15] V. Kac, *"Infinite Dimensional Lie Algebras"*, Birkhauser, 1983.
- [16] T. Fischbacher and H. Nicolai, *Low Level Representations for E₁₀ and E₁₁*, Contribution to the Proceedings of the Ramanujan International Symposium on Kac-Moody Algebras and Applications, ISKMAA-2002, Jan. 28-31, Chennai, India. hep-th/0301017.

- [17] N.D. Lambert, P.C. West, *Coset Symmetries in Dimensionally Reduced Bosonic String Theory*, Nucl.Phys. B615 (2001) 117-132, hep-th/0107209.
- [18] F. Englert, L. Houart, A. Taormina, P. West, *The Symmetry of M-Theories*, JHEP 0309 (2003) 020, hep-th/0304206.
- [19] A. Kleinschmidt and P. West, *Representations of G_{++} and the role of space-time*, JHEP 0402 (2004) 033, hep-th/0312247.